





Arradaza NV, Frobish D, et al. Reduced time for urinary alkalinization before high-dose methotrexate with preadmission oral bicarbonate. *J Oncol Pharm Pract* 2012;18:239-44.

[6] Von Hoff DD, Penta JS, Helman LJ, Slavik M. Incidence of drug-related deaths secondary to high-dose methotrexate and citrovorum factor administration. *Cancer Treat Rep* 1977;61:745-8.

[7] Sand TE, Jacobsen S. Effect of urine pH and flow on renal clearance of methotrexate. *Eur J Clin Pharmacol* 1981;19:453-6.

[8] Stoller RG, Hande KR, Jacobs SA, Rosenberg SA, Chabner BA. Use of plasma pharmacokinetics to predict and prevent methotrexate toxicity. *N Engl J Med* 1977;297:630-4.

[9] Chang JE, Medlin SC, Kahl BS, Longo WL, Williams EC, Lionberger J, et al. Augmented and standard Berlin-Frankfurt-Munster chemotherapy for treatment of adult acute lymphoblastic leukemia. *Leuk Lymphoma* 2008;49:2298-307.

[10] Khadela A, Vyas B, Joshi S. Role of glucarpidase for intoxication of high dose methotrexate toxicity. *J Pharm Appl Sci* 2019;6:110.

[11] Howard SC, McCormick J, Pui CH, Buddington RK, Harvey RD. Preventing and managing toxicities of high-dose methotrexate. *Oncologist* 2016;21:1471-82.

[12] Dolai TK, Panigrahi A, De R, Kumar M, Mandal P, Bagchi B, et al. Outcome of ALL-BFM (Berlin-Frankfurt-Münster) 90 Protocol in childhood acute lymphoblastic leukemia in a resource constraint setting. Washington, DC: American Society of Hematology; 2015.

[13] Ferdousi SA, Akhter A, Nahar K, Islam A. High dose methotrexate and leucovorin rescue therapy in childhood malignancies: Experience in resource-limited country. *Bangladesh J Child Health* 2017;41:15-23.

[14] Bendall LJ, Bradstock KF. G-CSF: From granulopoietic stimulant to bone marrow stem cell mobilizing agent. *Cytokine Growth Factor Rev* 2014;25:355-67.