

肝科门诊护理不良事件的原因及防范措施

周蓓虹

上海中医药大学附属龙华医院 上海 200032

【摘	要】:目的:				方法:		2021	2~8	2021	9	~20)22
2	50				2021	2~8				20	21	9
~20	22 2			结果:			6.00%			82.00)%	
	68.00%	100		18						2		
		P<0.05	结论:									

【关键词】:

Causes and Preventive Measures of Nursing Adverse Events in Liver Outpatient Department

Beihong Zhou

Longhua Hospital Shanghai University of Traditional Chinese Medicine Shanghai 200032

Abstract: Objective: To explore the causes and preventive measures of adverse events in outpatient nursing of liver department. Methods: A total of 50 liver patients admitted to the hospital from February to August 2021 and from September 2021 to February 2022 were selected for research, and they were divided into observation group and control group. The nursing quality control group was the control group, and the nursing quality control group established from September 2021 to February 2022 was the observation group. Results: The incidence of nursing adverse events in the observation group was only 6.00%, and its nursing satisfaction was as high as 82.00%, which was higher than that in the control group, which was 68.00%; a total of 18 adverse events occurred in 100 patients, of which the proportion of nursing staff was not standardized. Highest, poor communication second. Compared with the two groups, the difference was statistically significant (P<0.05). Conclusion: The quality of nursing care in hepatobiliary surgery directly affects the prognosis of patients. Nurses need to analyze the causes of adverse nursing events in detail, formulate more targeted preventive measures, strengthen supervision, and create a harmonious and harmonious relationship between nurses and patients.

Keywords: Hepatology; Outpatient Department; Nursing adverse events; Reason; Preventive measures

[3]

1 资料与方法 2

1.1 20021 2-8 2021 9 ~2022 2 50 28 22 25-65 43.17



3

4

5

1.2.2

1

3

4

1.3 1 2 20 5 100 60~80 >80 <60 2 1.4 SPSS22.0 t

P 0.05

2 结果

2.1

6.00%

P<0.05				
1			[n %]
	50	3	6.00%	
	50	15	30.00%	
X ²	1	-	21.962	
Р	-	-	0.001	

2.2 5 100 18

	2	[n	%]
6			
		n=18	
		9	50.00%
		5	27.78%
		2	11.12%
5		1	5.55%
		1	5.55%

5



2.3

	\sim	\sim
レィ	11	11

	3			[n	%]
	50	30 60.00%	11 22%	9 18.00%	41 82.00%
	50	15 30.00%	19 38.00%	16 32.00%	34 68.00%
X ²	-				6.194
Р	-				0.000

23

P<0.05

	3			[n	%]
	50	30 60.00%	11 22%	9 18.00%	41 82.00%
	50	15 30.00%	19 38.00%	16 32.00%	34 68.00%
X ²	-				6.194
Р	-				0.000

[5]

3 讨论

参考文献:

[1] . [J]. ,2020,17 (21):

3-6.

[2] , .

[J]. ,2021,6 (32):

3-5+9.

[3] , , , .

[7]

[J].

,2020,26(14):6.

[4] , , .

[J]. ,

2020, 49(001): 3-8.

[5] , .

[J]. ,

[8]

2019,34 (24):4-9.

[6] , .

[J].2021,7(89):189-

2

临床医学研究: 2022 年 4 卷 10 期

ISSN: 2705-0939(Print); 2705-0475 (Online)



192 [J]. : ,2021,19
[7] , , , , . (003):5-8.

[J]. , 2022, [9] , .

29(005):12-14.

[8] , . 316 130-132.